

NEW MEDICINE APPLICATIONS BROUGHT TO THE BORDERS FORMULARY COMMITTEE (BFC) AND AREA DRUG & THERAPEUTICS COMMITTEE (ADTC) 2016/17

New Medicine Product/Device Endorsement Categories

Α	Approved for general use - hospital and general practice
В	Approved for specialist use only
С	Approved for consultant initiation, with ongoing prescribing in general practice
D	Approved for shared care between hospital and general practice. Shared care protocol to be developed. (Information included here)
Е	Not approved (reason given)
F	No decision – further information required (listed here)

Green	Unrestricted General Use Used widely and in accordance with a respectable, responsible body of clinical opinion (e.g. BNF for children; SIGN/NICE recommendation; royal college guidelines)
Amber	General Use With Restrictions (Prescribers to be outlined) Supporting clinical evidence has been evaluated by NHS Borders ADTC/Formulary committee. Local use has peer group support and reflects practice in other NHS Board areas. May require "shared care protocol".(include information here) Specific patient consent not normally required.
Red	Specialist Use Only Supporting clinical evidence has been evaluated by NHS Borders ADTC/Formulary committee. Limited evidence of efficacy available. Rarely used or may have serious potential side effects requiring close supervision of treatment Specific patient consent indicated.
Black	Not Approved for use

NEW MEDICINE APPLICATIONS BROUGHT TO BFC MEETING OF 13 th APRIL 2016		
NMA	Indication	BFC
		Decision
Alendronate	Treatment of Osteoporosis – fracture prevention	Approved Category C –
(Binosto)		limited numbers -
		feedback/additional info
		required
Certolizumab	In combination with methotrexate for the treatment of moderate to severe active	Approved Category B
pegol (Cimzia)	rheumatoid arthritis in adult patients when the response to disease modifying anti-	
	rheumatic drugs, including methotrexate has been inadequate. As monotherapy in	

	case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate.	
Tolvaptan (Jinarc)	To slow the progression of cyst development and renal insufficiency of autosomal dominant polycystic kidney disease (ADPKD) in adults with chronic kidney disease stage 1 to 3 at initiation of treatment with evidence of rapidly progressing disease.	Approved Category B
Sacubitril/valsartan (Entresto)	Treatment of symptomatic chronic heart failure with reduced ejection fraction.	Approved Category C

NEW MEDICINE APPLICATIONS BROUGHT TO ADTC MEETING OF 11 th MAY 2016		
NMA	Indication	ADTC
		Decision
Tropicamide &	To obtain pre-operative mydriasis or for diagnostic purposes when monotherapy is	Approved Category B
Phenylephrine	known to be insufficient.	once evidence and info
(Mydriasert)		provided
Elvitegravir,	Treatment of adults and adolescents (aged 12 years and older with body weight at	Approved Category B
cobicistat,	least 35kg) infected with human immunodeficiency virus-1 (HIV-1) without any	
emtricitabine,	known mutations associated with resistance to the integrase inhibitor class,	
tenofovir	emtricitabine or tenofovir.	
alafenamide film		
coated tablet		
(Genvoya)		
Adalimumab	Treatment of active moderate to severe hidradenitis suppurativa (HS) (acne inversa)	Approved Category B
(Humira)	in adult patients with an inadequate response to conventional systemic HS therapy.	

NEW MEDICINE APPLICATIONS BROUGHT TO BFC MEETING OF 8 th JUNE 2016		
NMA	Indication	BFC
		Decision
Camellia sinensis green tea leaf extract	Cutaneous treatment of external genital and perianal warts (<i>condylomata acuminata</i>) in immunocompetent patients from the age of 18 years.	Approved Category B
(Catephen)		
Febuxostat	Prevention and treatment of hyperuricaemia in adult patients undergoing	Approved Category B
(Adenuric)	chemotherapy for haematologic malignancies at intermediate to high risk of Tumour Lysis Syndrome (TLS).	

NEW MEDICINE APPLICATIONS BROUGHT TO ADTC MEETING OF 13th JULY 2016

NMA	Indication	ADTC Decision
Sugammadex (Bridion)	Reversal of neuromuscular blockade induced by Rocuronium or Vecuronium.	Approved Category B
Secukinumab (Cosentyx)	Treatment of active ankylosing spondylitis (AS) in adults who have responded inadequately to conventional therapy.	Approved Category B

NEW MEDICINE APPLICATIONS BROUGHT TO BFC MEETING OF 10 th AUGUST 2016		
NMA	Indication	BFC Decision
Secukinumab (Cosentyx)	Alone or in combination with methotrexate, for the treatment of active psoriatic arthritis in adult patients when the response to previous disease-modifying anti-rheumatic drug (DMARD) therapy has been inadequate.	Approved – Category B
AquaMax Emollient Cream	Emollient, for dry and itchy skin conditions such as Eczema and Psoriasis.	Approved – Category A
Guanfacine prolonged-release (Intuniv)	Treatment of attention deficit hyperactivity disorder (ADHD) in children and adolescents 6 to 17 years old for whom stimulants are not suitable, not tolerated or have been shown to be ineffective. Treatment must be used as part of a comprehensive ADHD treatment programme, typically including psychological, educational and social measures.	Not Approved – Category E
Certolizumab pegol (Cimzia)	Treatment of adult patients with severe active axial spondyloarthritis, comprising: Ankylosing spondylitis (AS) Adults with severe active ankylosing spondylitis who have had an inadequate response to, or are intolerant to non-steroidal anti-inflammatory drugs (NSAIDs); Axial spondyloarthritis without radiographic evidence of AS (nr-axSpA). Adults with severe active axial spondyloarthritis without radiographic evidence of AS but with objective signs of inflammation by elevated CRP and /or MRI, who have had an inadequate response to, or are intolerant to NSAIDs	Approved – Category B
Certolizumab pegol (Cimzia)	In combination with methotrexate, for the treatment of active psoriatic arthritis in adults when the response to previous disease-modifying antirheumatic drug (DMARD) therapy has been inadequate. Certolizumab pegol can be given as monotherapy in case of intolerance to methotrexate or when continued treatment with methotrexate is inappropriate	Approved – Category B
Adalimumab (unlicensed indication)	Pyoderma gangrenosum	Approved – Category B for 1 patient

NEW MEDICINE APPLICATIONS BROUGHT TO ADTC MEETING OF 14 th SEPTEMBER 2016		
NMA	Indication	ADTC
		Decision
Calcipotriol +	Topical treatment of psoriasis vulgaris in adults.	Approved Category F
Betamethasone		
(Enstilar)		
Paliperidone	A three-monthly injection is indicated for the maintenance treatment of schizophrenia	Approved Category B
palmitate	in adult patients who are clinically unstable on one-monthly paliperidone palpitate	
(Trevicta)	injectable product.	
Idarucizumab	Reversal of dabigatran etexilate	Approved Category B
(Praxbind)		
OM-89	UTI Prophylaxis in female patients with recurrent multiple resistant, uncomplicated	Approved Amber up to 5
(UroVaxam)	UTI	patients
Vortioxetine	Treatment of major depressive episodes in adults.	Approved Category F
(Brintellix)		

NEW MI	NEW MEDICINE APPLICATIONS BROUGHT TO BFC MEETING OF 12 th OCTOBER 2016		
NMA	Indication	BFC	
		Decision	
Budesonide 9mg prolonged release tablet(Cortiment)	Induction of remission in patients with mild to moderate active ulcerative colitis (UC) where aminosalicylate (5-ASA) treatment is not sufficient.	Category F	
Aflibercept	Treatment of visual impairment due to myopic choroidal neovascularisation (myopic	Category B	
40mg/ml (Eylea)	CNV)		
Alirocumab (Praluent®)	Primary hypercholesterolaemia (heterozygous familial and non-familial) or mixed dyslipidaemia, as an adjunct to diet: - in combination with a statin or statin with other lipid lowering therapies in patients unable to reach LDL-C goals with the maximum tolerated dose of a statin or, alone or in combination with other lipid-lowering therapies in patients who are statin-intolerant, or for whom a statin is contraindicated.	Category B	

NEW MEDICINE APPLICATIONS BROUGHT TO ADTC MEETING OF 9 th NOVEMBER 2016		
NMA	Indication	ADTC
		Decision
No applications		

NEW MEDICINE APPLICATIONS BROUGHT TO BFC MEETING OF 14 th DECEMBER 2016		
NMA	Indication	BFC
		Decision
Aclidinium/Formoterol	COPD	Category A
(Duaklin Genuair)		
Insulin Degludec	Treatment of diabetes mellitus in adults.	Category F
(Tresiba)		
Dulaglutide (Trulicity)	Type 2 diabetes mellitus to improve glycaemic control as add-on therapy in	Category C
	combination with other glucose-lowering medicinal products including insulin,	
	when these, together with diet and exercise, do not provide adequate glycaemic	
	control.	
Melatonin MR	Acute insomnia in Critical Care.	Category B/Amber
Alemtuzumab	Adult patients with relapsing-remitting multiple sclerosis (RRMS) with active	Category B
(Lemtrada)	disease defined by clinical or imaging features.	
Sofosbuvir-Velpatasvir	Treatment of chronic hepatitis C virus (HCV) infection in adults.	Category B
(Epclusa)		
Tapentadol SR	Management of severe chronic pain in adults, which can be adequately managed	Category F
	only with opioid analgesics.	

NEW MEDICINE APPLICATIONS BROUGHT TO ADTC MEETING OF 11 th JANUARY 2017				
NMA	Indication	ADTC		
		Decision		
No applications				

NEW MEDICINE APPLICATIONS BROUGHT TO BFC MEETING OF 8 th FEBRUARY 2017				
NMA	Indication	BFC		
		Decision		
Lurasidone	Treatment of schizophrenia in adults aged 18 years and over	Category C		
(Latuda)				
Elbasvir 50mg,	Treatment of chronic hepatitis C (CHC) in adults. (The efficacy of elbasvir-	Category B		
Grazoprevir	grazoprevir has not been demonstrated in genotypes 2, 3, 5 and 6. Elbasvir-			
100mg film-	grazoprevir is not recommended in patients infected with these genotypes).			
coated tablet				
(Zepatier)				

Evolocumab 140mg (Repatha)	Primary hypercholesterolaemia (heterozygous familial hypercholesterolaemia and non-familial) or mixed dyslipidaemia, as an adjunct to diet: • in combination with a statin or statin with other lipid lowering therapies in patients unable to reach low density lipoprotein-cholesterol (LDL-C) goals with the maximum tolerated dose of a statin or, • alone or in combination with other lipid-lowering therapies in patients who are statin-intolerant, or for whom a statin is contraindicated. See SMC Advice for restrictions	Category B
5-Fluorouracil for	Extensive Keloid Scarring	Category B – non
injection		formulary off label request

NEW MEDICINE APPLICATIONS BROUGHT TO ADTC MEETING OF 8 th MARCH 2017				
NMA	Indication	ADTC		
		Decision		
Tocilizumab	In combination with methotrexate is indicated for the treatment of juvenile idiopathic	Category B – 1 patient		
(RocActemra)	polyarthritis (rheumatoid factor positive or negative and extended oligoarthritis) in			
	patients 2 years of age and older, who have responded inadequately to previous			
	therapy with methotrexate.			
Aztreonam	Alternative to carbapenems in treatment of severe infection caused by susceptible	Category B		
(Azactam)	gram negative bacteria on the advice of Consultant Microbiologist			